



# Central and North Florida Regional Turf Spray Program

## Non-Dormant Bermudagrass – Warm Season

With this geography-specific spray program, superintendents have a customized rotation management plan for disease control. Be sure and include these important rotations in your plan and have strong, resilient turf all year long.

- Intrinsic® brand fungicides provide a strong foundation of plant health benefits, including broad-spectrum disease control, stronger roots and higher stress tolerance.
- The turf-safe Maxtima® fungicide is a DMI that delivers long-lasting disease control of key diseases at any temperature for rotation flexibility and year-round confidence.
- The broad-spectrum DMI power of Navicon® Intrinsic® brand fungicide controls more than 20 major diseases while improving overall turf health from root to blade.

Learn how to create impressive playing conditions on your course at [betterturf.basf.com](http://betterturf.basf.com)





# Central and North Florida Non-Dormant Bermudagrass

Long, cool springs can affect turfgrass growth in this region, though spring and fall tend to remain temperate. While turfgrass does not go dormant in winter, temperatures can dip below the 55-degree growing temperature for bermudagrass. Winter is considered the dry season, but unexpected weeks of clouds and rain sometimes occur. Summer months are humid, and the tropical season in late July and August can be wet and very cloudy for weeks at a time.

14-28  
day program



Timing	Product	Active Ingredient	Rate/1,000 sq. ft.	Key Diseases & Benefits
1 Aug	<b>Encartis™ fungicide</b>	boscalid + chlorothalonil	4.0 fl oz	Algae, leaf spots, tropical weather protection
	Fore® 80WP Rainshield® specialty fungicide	mancozeb	SLR <sup>1</sup> (Standard Local Rate)	
2 Sept	<b>Navicon® Intrinsic® brand fungicide</b>	pyraclostrobin + mefentrifluconazole	0.85 fl oz (water-in) <sup>2</sup>	Leaf spots, spring dead spot, take-all root rot, tropical weather protection
	Signature™ XTRA Stressgard® fungicide + Daconil® WeatherStik® fungicide	fosetyl-Al (aluminum-tris) + chlorothalonil	SLR	
3 Oct	<b>Navicon Intrinsic brand fungicide</b>	pyraclostrobin + mefentrifluconazole	0.85 fl oz (water-in)	Spring dead spot, take-all root rot
	Signature XTRA Stressgard fungicide + Daconil WeatherStik fungicide	fosetyl-Al (aluminum-tris) + chlorothalonil	SLR	
4 Nov	<b>Lexicon® Intrinsic brand fungicide</b>	pyraclostrobin + fluxapyroxad	0.47 fl oz (water-in)	Spring dead spot, take-all root rot, winter protection
5 Dec	<b>Encartis fungicide</b>	boscalid + chlorothalonil	4.0 fl oz	Algae, leaf spots, Microdochium Patch <i>Pythium</i> (if needed)
	Segway® fungicide	cyazofamid	SLR	
6 Jan	Signature XTRA Stressgard fungicide + 26GT fungicide	fosetyl-Al (aluminum-tris) + iprodione	SLR	Algae, leaf spots, Microdochium Patch <i>Pythium</i> (if needed)
	Segway fungicide	cyazofamid	SLR	
7 Feb	<b>Navicon Intrinsic brand fungicide</b>	pyraclostrobin + mefentrifluconazole	0.85 fl oz (water-in)	Fairy ring, leaf spots, Microdochium patch, take-all root rot
	26GT fungicide	iprodione	SLR	
8 Mar	<b>Lexicon Intrinsic brand fungicide</b>	pyraclostrobin + fluxapyroxad	0.47 fl oz (water-in)	Fairy ring, leaf spots, take-all root rot
9 Apr	Mirage® Stressgard fungicide	tebuconazole	SLR (water-in)	Fairy ring, leaf spots, take-all root rot
10 May	<b>Xzemplar® fungicide</b>	fluxapyroxad	0.26 fl oz (water-in)	Fairy ring, leaf spots, mini ring
	Signature XTRA Stressgard fungicide + Daconil WeatherStik fungicide	fosetyl-Al (aluminum-tris) + chlorothalonil	SLR	
11 Jun	<b>Lexicon Intrinsic brand fungicide</b>	pyraclostrobin + fluxapyroxad	0.47 fl oz (water-in)	Fairy ring, leaf spots, mini ring, summer stress
12 Jul	<b>Xzemplar fungicide</b>	fluxapyroxad	0.26 fl oz (water-in)	Fairy ring, leaf spots, mini ring
	Medallion® fungicide + Daconil WeatherStik fungicide	fludioxonil + chlorothalonil	SLR	

Multiple products listed per month are meant to be a rotational option to control additional disease pressure that may occur.

<sup>1</sup> Standard local rate for the products shown; consult label and/or local extension for regional recommendations

<sup>2</sup> Recommended application spray volume is 2-4 gallons/1000ft<sup>2</sup>. Provide 1/4" irrigation post application. To improve hydrophobic conditions and improve fungicide efficacy, BASF recommends regular soil wetting agent applications during the growing season.

Product timing is a suggestion only; optimize application for local weather and field conditions. Confirm active status of product registration in your state before using; not all products are registered in all states. Application Rate/1,000 sq. ft. is based on maximum label rate or range. Claims related to root strength, stress tolerance, resiliency and growth efficiency are specific to those BASF products containing pyraclostrobin, as shown.

Always read and follow label directions.

Encartis is a trademark, and Intrinsic, Lexicon, Navicon and Xzemplar are registered trademarks of BASF. Signature is a trademark, and 26GT, Mirage and Stressgard are registered trademarks of Bayer Environmental Science. Fore and Rainshield are registered trademarks of Corteva. Daconil, Medallion and Weatherstik are registered trademarks of Syngenta Professional Products. Segway is a registered trademark of PBI Gordon.

© 2021 BASF Corporation. All Rights Reserved.

RGSP-S17-08/21