

14-28 day program

Northern California

Northern California

This area features moderate temperatures and variable rainfall, with spring and fall being the wettest seasons. Disease pressure is highest in coastal and valley areas where humidity and nighttime temperatures provide ideal conditions for diseases like brown ring patch (formerly known as *Waitea*) and summer patch. Fairy ring, anthracnose, snow mold and rapid blight are also common on this region's cool season greens, tees and fairways.

	Timing	Product	Active Ingredient	Rate/1,000 sq ft	Target Diseases & Benefits
1	Mar 1 ¹	Xzemplar® fungicide + Affirm™ WDG	Fluxapyroxad + Polyoxin-D	0.26 fl oz + 0.9 fl oz	Patch diseases
2	Apr 1	Lexicon® Intrinsic® brand fungicide + Fore® 80WP Rainshield®	Pyraclostrobin + Fluxapyroxad + Mancozeb	0.47 fl oz + 6.0-8.0 fl oz	Bermudagrass decline, Leaf spot, Brown ring patch, Take-all patch, Plant health
3	May 1	Navicon® Intrinsic® brand fungicide	Mefentrifluconazole + Pyraclostrobin	0.85 fl oz	Brown ring patch, Leaf spot, Plant Health
4	Jun 1 (Memorial Day)	Lexicon Intrinsic brand fungicide + Fore 80WP Rainshield	Pyraclostrobin + Fluxapyroxad + Mancozeb	0.47 fl oz + 6.0-8.0 fl oz	Brown ring patch, Summer patch, Fairy ring², Rapid blight, Anthracnose, Plant Health
5	Jul 1	Xzemplar fungicide + Daconil®	Fluxapyroxad + Chlorothalonil	0.26 fl oz + 3.0 fl oz	Fairy ring ² , Anthracnose
6	Jul 15 (Fourth of July)	Lexicon Intrinsic brand fungicide	Pyraclostrobin + Fluxapyroxad	0.47 fl oz	Anthracnose, Summer Plant Health
7	Aug 1	Segway®	Cyazofamid	0.9 fl oz	Pythium
8	Sep 1	Maxtima fungicide + Signature™ XTRA Stressgard®	Mefentrifluconazole + Fosetyl-AL (Aluminum tris)	0.8 fl oz + 4.0 fl oz	Anthracnose, Pythium
9	Oct 1 (Labor Day)	Lexicon Intrinsic brand fungicide	Pyraclostrobin + Fluxapyroxad	0.47 fl oz	Take-all patch, Bermudagrass decline, Spring dead spot, Plant Health
10	Oct 15	Trinity® fungicide + 26GT® + Medallion®	Triticonazole + Iprodione + Fludioxonil	1.0 fl oz + 4.0 fl oz + 0.5 fl oz	Snow mold

¹ Adjust first application date to reflect the onset of temperatures above 60° F.

Product timing is a suggestion only; optimize application for local weather and field conditions. Confirm active status of product registration in your state before using; not all products are registered in all states. Application Rate/1,000 sq. ft. is based on maximum label rate or range. Spray program for greens is designed to cover four acres. Plant health claims are related to root strength, stress tolerance, resiliency and growth efficiency and are specific to those BASF products containing pyraclostrobin, as shown.

Always read and follow label directions.

Intrinsic, Lexicon, Maxtima, Navicon, Trinity and Xzemplar are registered trademarks of BASF. Signature is a trademark and 26GT and Stressgard are registered trademarks of Bayer. Affirm is a trademark of Cleary Chemical LLC. Fore and Rainshield are registered trademarks of Dow AgroSciences. Segway is a registered trademark of ISK, Ltd. Daconil and Medallion are registered trademarks of Syngenta Professional Products.

² To improve hydrophobic conditions, make application of preferred wetting agent 1-2 weeks prior to first fungicide treatments for fairy ring.