

day program

Inland Northwest

Eastern WA, Eastern OR, Northern Idaho, Western Montana, Western Wyoming, and Northern Utah

This area features moderate temperatures and variable rainfall, with fall and winter being the wettest seasons. Disease pressure is highest in coastal and valley areas where irrigated turf and low mowing heights periodically provide ideal conditions for disease development. Leaf spots, patch diseases, fairy ring and snow mold are common on this region's cool-season greens, tees and fairways.

	Timing	Product	Active Ingredient	Rate/1,000 sq ft	Target Diseases & Benefits
1	Apr 1 ¹	Lexicon® Intrinsic® brand fungicide	Pyraclostrobin + Fluxapyroxad	0.47 fl oz	Rust, Leaf spot, Snow mold, Fairy ring ²
2	May 1	Navicon® Intrinsic® brand fungicide + Affirm™ WDG	Mefentrifluconazole + Pyraclostrobin + Polyoxin-D	0.85 fl oz + 0.9 fl oz	Leaf Spot, Brown ring patch, Necrotic ring spot, Brown patch
3	Jun 1 (Memorial Day)	Lexicon Intrinsic brand fungicide	Pyraclostrobin + Fluxapyroxad	0.47 fl oz	Broad Spectrum Disease Protection, Summer Plant Health
4	Aug 1	Maxtima® fungicide + Signature™ XTRA Stressgard®	Mefentrifluconazole + Fosetyl-AL (Aluminum tris)	0.8 fl oz + 4.0 fl oz	Anthracnose, Plant Health
5	Oct 1	Xzemplar® fungicide	Fluxapyroxad	0.26 fl oz	Fairy ring ²
8	Oct 15	Insignia® SC Intrinsic brand fungicide + Trinity® fungicide + Daconil®	Pyraclostrobin + Triticonazole + Chlorothalonil	0.7 fl oz + 1.0 fl oz + 3.0 fl oz	Snow mold, Winter protection

¹ Adjust first application date to reflect the onset of temperatures above 60° F.

Product timing is a suggestion only; optimize application for local weather and field conditions. Confirm active status of product registration in your state before using; not all products are registered in all states. Application Rate/1,000 sq. ft. is based on maximum label rate or range. Spray program for greens is designed to cover four acres. Plant health claims are related to root strength, stress tolerance, resiliency and growth efficiency and are specific to those BASF products containing pyraclostrobin, as shown.

Always read and follow label directions.

² To improve hydrophobic conditions, make application of preferred wetting agent 1-2 weeks prior to first fungicide treatments for fairy ring.